

Mouse Monoclonal Antibody to

Phosphotyrosine

clone 16F4

Order No.: 0030-100/PTYR-16F4

Size (µg) 100

Lot No.: 0030S



www.nanotools.de

orders & support:

email: info@nanotools.de

phone: +49-7641-455 670

fax: +49-7641-455 671

02/060307F

| Isotype | Species Reactivity | Applications | Mol. Weight | Ref.Cell Line | Epitope | Immunogen |
|---------|------------------------|---------------|-------------|---------------|----------------------------|----------------------------------|
| IgG1 | human, mouse, rat, dog | WB, ELISA, IP | pattern | | ... D - I - pY - A - E ... | phosphopeptide conjugated to KLH |

Background and Specificity:

Phosphorylation and dephosphorylation of cellular proteins are central steps in transducing extracellular signals to the cell nucleus. Phosphorylated epitopes may serve as docking sites for the assembly of protein complexes or may alter the 3-dimensional protein structure thus modulating enzymatic activity or the ability to undergo protein-protein-interactions. Modification of proteins on tyrosine residues is mediated by protein tyrosine kinases. Tyrosine phosphorylation may alter the biological activity or mediate the assembly of protein complexes via interaction of phosphotyrosine residues with SH2 or PID domains.

Mab PTYR-16F4 recognizes phosphotyrosine in the context of the surrounding amino acids, tolerating hydrophobic amino acids directly neighboured to phosphotyrosine.

Related Products

mab against Phosphotyrosine

#0027-100/pTyr-2C8

#0028-100/pTyr-3B12

#0029-100/pTyr-9H8

#0133-100/pTyr-2A5

#0156-100/pTyr-9F1

#0226-100/pTyr-1F9

Purification: The antibody was purified from serum-free cell culture supernatant by subsequent thiophilic adsorption and size exclusion chromatography.

Formulation: lyophilized from 1 ml 2 x PBS / 0.09 % Na-azide / PEG and Sucrose.

Reconstitution: Reconstitute with 1 ml H₂O (15 min, RT).

Stability: For long-term storage, freeze lyophilizate upon arrival (-20°C). Upon reconstitution, aliquote and freeze in liquid nitrogen; reconstituted antibody can be stored frozen at -80°C up to 1 year. Thaw aliquots at 37°C. Thawed aliquots may be stored at 4°C up to 3 months.

Avoid repeated freeze / thaw cycles.

Positive Control: #0038: phosphotyrosine MW standard


Immunoblotting: 0.5 µg/ml for HRPO/ECL detection
Recommended blocking buffer: Casein/Tween 20 based blocking and blot incubation buffer, e.g. nanoTools product #3031-500/CPPT or #3031-3000/CPPT.

Immunoprecipitation: use at 1 - 10 µg per 10⁶ pervanadate-treated A431 cells

Immunocytochemistry: ND

ELISA: use at 0.05 µg/ml

All products are supplied for research and investigational use only. Not for use in humans or laboratory animals.



Phosphotyrosine Detection

Lysates of pervanadate-treated A431 cells were probed with
 lane 1: mab 2A5 (IgG), 1 µg/ml
 lane 2: mab 2C8 (IgG), 1 µg/ml
 lane 3: mab 3B12 (IgG), 1 µg/ml
 lane 4: mab 9H8 (IgG), 1 µg/ml
 lane 5: mab 16F4 (IgG), 1 µg/ml
 lane 6: mab 9F1 (IgG), 1 µg/ml