

Mouse Monoclonal Antibody to

α -Catenin

clone 15D9

Order No.: 0001-100/a-CAT-15D9

Size (μ g) 100

Lot No.: 0001S



www.nanotools.de

orders & support:

email: info@nanotools.de

phone: +49-7641-455 670

fax: +49-7641-455 671

02/140307F

Isotype	Species Reactivity	Applications	Mol. Weight	Ref.Cell Line	Epitope	Immunogen
IgG1	human, mouse, rat, dog	WB, ELISA, IP, ICC	102 kDa	A431		recombinant α -catenin

Background and Specificity:

The α -, β - and γ -catenins are cytoplasmic proteins mediating the interaction of Ca^{2+} -dependent transmembrane adhesion molecules (cadherins) with the cytoskeletal network. The N-terminus of α -catenin directly interacts with exon 3 of β -catenin. The C-terminus of α -catenin interacts with the actin filament network.

Mab a-CAT-15D9 specifically recognizes α -catenin in Western blot at 102 kDa and is also suitable for ELISA, immunoprecipitation and immunocytochemistry.

Related Products

mab to b-catenin (N-Term/Exon2)

#0003-100/b-CAT-7D11

mab to b-catenin (Exon3)

#0004-100/b-CAT-9G2

mab to b-catenin (Core)

#0005-100/b-CAT-9G10

mab to b-catenin (C-Term/Exon14)

#0002-100/b-CAT-7D8

mab to b-catenin (C-Term)

#0006-100/b-CAT-10H8

mab to dephospho-b-catenin (aa35-50)

#0051-100/b-CAT-7A7

mab to dephospho-b-catenin (aa27-37)

#0052-100/b-CAT-8E4

mab to phospho-b-catenin (pY86)

#0123-100/b-CAT-24E1

mab to phospho-b-catenin (pY654)

#0159-100/b-CAT-1B11

For monoclonal antibodies against LEF, TFF3, and E-, M- and N-Cadherin, please refer to our website at www.nanotools.de

Purification:	The antibody was purified from serum-free cell culture supernatant by subsequent thiophilic adsorption and size exclusion chromatography.
Formulation:	lyophilized from 1 ml 2x PBS / 0.09 % Na-azide / PEG and Sucrose.
Reconstitution:	Reconstitute with 1 ml H_2O (15 min, RT).
Stability:	For long-term storage, freeze lyophilizate upon arrival (-20°C). Upon reconstitution, aliquote and freeze in liquid nitrogen; reconstituted antibody can be stored frozen at -80°C up to 1 year. Thaw aliquots at 37°C . Thawed aliquots may be stored at 4°C up to 3 months.

Avoid repeated freeze / thaw cycles.


Positive Control: #0831: Cell lysate from untreated A431 cells.

Immunoblotting: 0.5 $\mu\text{g}/\text{ml}$ for HRPO/ECL detection
Recommended blocking buffer: Casein/Tween 20 based blocking and blot incubation buffer, e.g. nanoTools product #3031-500/CPPT or #3031-3000/CPPT.

Immunoprecipitation: use at 1- 10 μg per 10^6 vanadate-treated A431 cells

Immunocytochemistry: use at 0.1-1 $\mu\text{g}/\text{ml}$; mab a-CAT-15D9 may tolerate formaldehyd

ELISA: use at 0.05 $\mu\text{g}/\text{ml}$



Detection of endogenous alpha catenin

Whole cell lysates of serum starved tumor cells (20.000 cells per lane) were applied to SDS-PAGE and transferred to a PVDF membrane. The immunoblot was probed with mab a-Cat-15D9 (0.5 $\mu\text{g}/\text{ml}$) for 1h at RT and developed by ECL (exp. time: 30 sec).

lane 1: A431; lane 2: A549; lane 3: SKOV3; l OVCA5; lane 5: HaCaT; lane 6: PC3; lane HepG2

All products are supplied for research and investigational use only. Not for use in humans or laboratory animals.