

Mouse Monoclonal Antibody to

Casein Kinase 1 epsilon (CK1ε)

clone 4D7

Order No.: 0172-100/CK1e-4D7

Size (µg) 100

Lot No.: 0172S



www.nanotools.de

orders & support:

email: info@nanotools.de

phone: +49-7641-455 670

fax: +49-7641-455 671

02/160307F

Isotype	Species Reactivity	Applications	Mol. Weight	Ref.Cell Line	Epitope	Immunogen
IgG1	human, mouse, rat, dog	WB, ELISA	45 kDa	A431	aa 355 - 380	peptide conjugated to hemocyanin

Background and Specificity:

Casein kinases are operationally defined by their preferential utilization of acidic proteins such as caseins as substrates. It can also phosphorylate a large number of proteins and participates in Wnt signaling.

Mab CK1e-4D7 specifically recognizes the epsilon isoform of CK1.

Related Products

mab to b-catenin (N-Term/Exon2)

#0003-100/b-CAT-7D11

mab to b-catenin (Exon3)

#0004-100/b-CAT-9G2

mab to b-catenin (Core)

#0005-100/b-CAT-9G10

mab to b-catenin (C-Term/Exon14)

#0002-100/b-CAT-7D8

mab to b-catenin (C-Term)

#0006-100/b-CAT-10H8

mab to dephospho-b-catenin (aa35-50),

#0051-100/b-CAT-7A7

mab to dephospho-b-catenin (aa27-37),

#0052-100/b-CAT-8E4

mab to phospho-b-catenin (pY86)

#0123-100/b-CAT-24E1

mab to phospho-b-catenin (pY654)

#0159-100/b-CAT-1B11

For monoclonal antibodies against alpha-catenin, LEF, TFF3, and M-Cadherin, please refer to our website at www.nanotools.de

Purification:	The antibody was purified from serum-free cell culture supernatant by subsequent thiophilic adsorption and size exclusion chromatography.
Formulation:	lyophilized from 1 ml 2 x PBS / 0.09 % Na-azide / PEG and Sucrose.
Reconstitution:	Reconstitute with 1 ml H ₂ O (15 min, RT).
Stability:	For long term storage, freeze lyophilizate upon arrival (-20°C). Upon reconstitution aliquote and freeze in liquid nitrogen; reconstituted antibody can be stored frozen at -80°C up to 1 year. Thaw aliquots at 37°C. Thawed aliquots may be stored at 4°C up to 3 months.


Avoid repeated freeze/thaw cycles.

Positive Control:	#0831: Cell lysate from untreated A431 cells
Immunoblotting:	0.5 µg/ml for HRPO/ECL detection Recommended blocking buffer: Casein/Tween 20 based blocking and blot incubation buffer, e.g. nanoTools product #3031-500/CPPT or #3031-3000/CPPT.

Immunoprecipitation: ND

Immunocytochemistry: ND

ELISA: use at 0.1 µg/ml



Detection of endogenous Casein Kinase 1 epsilon
Whole cell lysates of serum starved tumor cells (20.000 cells per lane) were applied to SDS-PAGE and transferred to a PVDF membrane. The immunoblot was probed with mab CK1e-4D7 (0.5 µg/ml) for 1h at RT and developed by ECL (exp. time: 30 sec).
lane 1: HeLa; lane 2: HepG2; lane 3: HEK293; lane 4: SH-SY5Y; lane 5: MDCK; lane 6: PC12; lane 7: CMT 93; lane 8: Neuro 2A; lane 9: NIH 3T3

All products are supplied for research and investigational use only. Not for use in humans or laboratory animals.